arXiv:nlin/0206042v1 [nlin.CG] 26 Jun 2002

Headway distribution of the asymmetric simple exclusion model

Milan Krbalek[a,b]
[a] Institute of Physics, Czech Academy of Science,
Cukrovarnická 10, Prague, Czech Republic
[b] Faculty of Nuclear Sciences and Physical Engineering,
Trojanova 13, Prague, Czech Republic
e-mail: milan.krbalek@uhk.cz

February 8, 2008

Abstract

We present an exact solution of headway distribution of the asymmetric simple exclusion model with open boundary conditions and compare it to the headway distributions of the highway traffic.

1 Introduction

Recently, a new tendency has appeared in the physics of traffic systems. Besides macroscopic characteristics of traffic (flow, density, etc.), researchers are focused on microscopic properties as well ([1]). One of these properties is the distribution of the distances (or time intervals) between neighboring cars, so called headway distribution. This function was investigated for existing traffic models and compared to the various regimes of the real traffic. It was observed that the real traffic data as well as many traffic models lead to a headway distribution which resembles the spacing distribution of the random-matrices eigenvalues ([2],[3],[8]). To verify this observation, we investigate the asymmetric simple exclusion process (ASEP) which is the well known exactly solvable model of traffic ([4],[5],[6]). We find the exact formula for headway distribution of the asymmetric simple exclusion model and compare it with the eigenvalues distributions obtained from random matrix theory.

In the traffic theory, the spacing distribution which was often called *headway distribution* $P(s)$ is defined as probability distribution of the distances s between neighboring cars (particles of the model). For the sake of simplicity we assume that $\langle s\rangle \equiv \int_0^\infty s\,P(s)\,ds = 1$. Among many others, we discern two special traffic systems in real roads. The first is the *bus transport* where no fixed schedule exists (see [2],[8]) and where the interaction between bus drivers exists owing to the check points in which the information about the time interval to the previous bus is transmitted to a driver. It was found that the headway distribution of that bus transport behaves as

$$P(s) = A\,s^2\,e^{-Bs^2}, \tag{1}$$

where A, B are constants determined by conditions $\int_0^\infty P(s)\,ds = 1$ and $\int_0^\infty s\,P(s)\,ds = 1$. The equation (1) originates from random matrix theory where it describes the distribution of

distances between eigenvalues of the unitary random matrices.
The second system described here is *highway traffic* for which we distinguish three basic regimes ([2]):

- a) *free-flow regime*, where the density of cars (number of cars per unit of length) is low and the cars are therefore moving as free. Headway of this regime is reduced to the Poissonian distribution

$$P_0(s) = e^{-s} \tag{2}$$

- b) *synchronized regime*, where the mean velocity is considerably reduced compared to the free-flow states due to a higher density of cars, but cars are still moving (no traffic jam). Headway distribution of these states is well described by the formula

$$P_\nu(s) = \frac{(\nu+1)^{\nu+1}}{\Gamma(\nu+1)} s^\nu e^{-(\nu+1)\,s}, \tag{3}$$

  where ν is a free parameter and Γ is Gamma function.

- c) *stop-and-go regime*, where cars are moving in traffic jams

Note that it is possible to obtain the formula (2) taking the limit $\nu \to 0^+$ in formula (3). Distribution $P_1(s)$ used to be called as semi-Poissonian statistics.

Roughly, particles in the free-flow regime do not interact between each other, contrary to the synchronized states when the interaction leads to the particle repulsion and to the fact that $\lim_{s\to 0^+} P_\nu(s) = 0$ for positive ν. One of aims of this article is to answer whether it is possible to get a synchronized state for the asymmetric simple exclusion model by an appropriate choice of input parameters, or the model is in the free regime for all parameters.

2 ASEP with open boundaries

Consider sites located on a chain of length N. Each site i $(1 \le i \le N)$ is either occupied by a particle, or empty. During the time interval dt, each particle jumps to the next site (in the defined direction) with probability dt if the target site is empty and does not hop otherwise. Moreover, during the time interval dt, a particle can enter the first chain position ($i = 1$) with probability $\alpha\, dt$ ($\alpha \in \langle 0, 1\rangle$) if this position is empty and a particle at the site $i = N$ can leave the chain with probability $\beta\, dt$ ($\beta \in \langle 0, 1\rangle$) if this site is occupied. These rules define a simple traffic model with a hard-core repulsion which decreases the probability for occurrence of the two particles close each other. This was *de facto* the reason why we have chosen ASEP for our investigation.
Two alternative methods of exact solution of the model are outlined in [4]. We will use the *matrix method* in our computations. Consider the matrices D, E and vectors $\langle w|$ and $|v\rangle$ satisfying the following algebraic rules

$$DE = D + E \tag{4}$$

$$\langle w|E = \frac{1}{\alpha}\langle w| \tag{5}$$

$$D|v\rangle = \frac{1}{\beta}|v\rangle \tag{6}$$

Then we can determine the probability of the arbitrary configuration $\mathcal{C} = (n_1, m_1, n_2, m_2, \ldots)$ of the model through the formula

$$P(\mathcal{C}) = \frac{1}{Z_N}\,\frac{\langle w|D^{n_1}E^{m_1}D^{n_2}E^{m_2}\ldots|v\rangle}{\langle w|v\rangle}, \tag{7}$$

where n_1 is the number of the empty sites, starting with the entering site of the chain (and (n_1+1)-th site is occupied), m_1 is the number of occupied sites starting with the site number n_1+1 (and (n_1+m_1+1)-th site is empty), and so on. Z_N is a normalization constant. Hence $\sum_i n_i + \sum_j m_j = N$. Owing to the fact

$$(D+E)^n = \sum_{m=1}^{n} \frac{m\,(2n-m-1)!}{n!(n-m)!} \sum_{i=0}^{m} E^i D^{m-i} \tag{8}$$

we can rewrite the normalization factor Z_N to the form

$$Z_N \equiv \frac{\langle w|(D+E)^N|v\rangle}{\langle w|v\rangle} = \sum_{m=1}^{N} \frac{m\,(2N-m-1)!}{N!(N-m)!} \sum_{i=0}^{m} \left(\frac{1}{\alpha}\right)^i \left(\frac{1}{\beta}\right)^{m-i} \tag{9}$$

Hence,

$$Z_N = \sum_{m=1}^{N} \frac{m\,(2N-m-1)!}{N!(N-m)!}\,\frac{\beta^{-m-1}-\alpha^{-n-1}}{\beta^{-1}-\alpha^{-1}} \tag{10}$$

These are the known results [4] which will be used below.

3 Headway distribution in the ASEP (a special case)

Our aim is an analytical expression for the spacing distribution, which means the probability of the gap of the length k behind the particle being on the i-th site. One can determine this probability $P^{(i)}(k)$ from the equation (7) as follows

$$P^{(i)}(k) = \frac{\mathcal{A}}{Z_N\,\langle w|v\rangle}\,\langle w|(D+E)^{i-1}D\,E^{k-1}D\,(D+E)^{N-i-k}\,|v\rangle, \tag{11}$$

since the first $i-1$ sites could be arbitrary occupied or not, i-th site is occupied, the following $k-1$ sites are empty, the site $i+k$ is occupied and the rest sites could be arbitrary occupied or not. If these rules are satisfied, one can see the gap of the length k behind the i-th position. The constant $\mathcal{A}$ can be calculated from the condition

$$\sum_{k=1}^{N-i} P^{(i)}(k) = 1$$

Generally, the matrices D, E are infinite-dimensional, but provided the condition

$$\alpha + \beta = 1 \tag{12}$$

one can find D, E to be real numbers. The equations (5) and (6) imply that $D = 1/\beta, E = 1/\alpha$ and the rule (4) imposes that ${}^{1}/_{\alpha} + {}^{1}/_{\beta} = {}^{1}/_{\alpha\beta}$, which leads to the equation (12). Instituting (12) and (9) into the (11) we obtain

$$P^{(i)}(k) = \mathcal{A}\,\alpha^2\,\beta^{k-1}\,(k \geq 1) \tag{13}$$

Strictly speaking, the derivation of the formula (13) from the equation (11) is possible only for $k \geq 3$ since for $k = 1, 2$ the relation (11) has a different form. But it can easily be verified that (13) holds true also for $k = 1, 2$. Note now that the result (13) is independent of the value i. For large N, after the re-scaling to $\langle s \rangle = 1$, equation (13) changes to the form $P(s) := P(k-1) = e^{-s}$, which is the Poissonian distribution. This clearly shows that under the condition (12), the interaction among the particles vanishes in the continuous limit and hence the movement of the particles is asymptotically free (in the sense mentioned in the section 2). Also the real headway distribution (without the re-scaling) can be determined in this section. If (12) holds true then

$$P^{(i)}(k) = \mathcal{N}\left(\frac{1}{\alpha}\right)^{N-2}\left(\frac{1}{\beta}\right)^{N+1-k}, \tag{14}$$

when the normalization pre-factor $\mathcal{N}$ can be computed either from the condition $\sum_{k=1}^{N-i} P^{(i)}(k) = 1$ (through all the lengths of the gap) or from the equation

$$\frac{1}{\mathcal{N}} = \frac{\langle w|(D+E)^{N-i}D\,(D+E)^{N-i}|v\rangle \;-\; \langle w|(D+E)^{i-1}D\,E^{N-i}|v\rangle}{\langle w|v\rangle},$$

which takes into account all the possible combinations with occupied site number i and rejects the combinations when all the sites (behind the site number i) are empty. This provides

$$\frac{1}{\mathcal{N}} = \left(\frac{1}{\alpha}\right)^{N-i}\left(\frac{1}{\beta}\right)^{N}(1-\beta^{N-i}),$$

which together with (14) yields

$$P^{(i)}(k) = \frac{\alpha}{1-\beta^{N-i}}\,\beta^{k-1}$$

In the large N limit, one can calculate from (12) that

$$P^{(i)}(k) = \alpha\,\beta^{k-1}$$

and hence

$$\langle k \rangle \equiv \sum_{k=1}^{\infty} k\,P^{(i)}(k) = \alpha \sum_{k=1}^{\infty} k\,\beta^{k-1} = \frac{1}{\alpha}$$

for the mean headway.

4 Headway distribution of the ASEP (general case)

In this section we are solving the ASEP (equation (11), strictly speaking) for arbitrary values of the parameters α, β. Introduce now the following pre-factors

$$\omega = \omega(m,k,i,N,p) = \frac{m\,p\,(2N-2i-2k-m-1)!(2i-p-3)!}{(N-i-k)!(N-i-k-m)!(i-1)!(i-p-1)!}$$

$$A_1 = A_1(p,q,a,w) = \left(\begin{array}{c} p-q-a+w-1 \\ p-q-1 \end{array} \right)$$

$$A_2 = A_2(p,q,b,w) = \left(\begin{array}{c} p-q-b+w-1 \\ w-1 \end{array} \right)$$

With the help of them and the computations presented in an Appendix we obtain the desired result



FIGURE 1. Headway distribution of the ASEP . The curve is (2). Bars and plus signs show the result (15) after the re-scaling to mean headway equal to one for $\alpha = 0.15, \beta = 0.6$ and $\alpha = 0.3, \beta = 0.5$, respectively.

$$P^{(i)}(k) = \frac{\mathcal{A}}{Z_N} \sum_{p=1}^{i-1} \sum_{m=1}^{N-i-k} \sum_{l=0}^{m} \omega \left(S_1 + S_2 + S_3 + S_4 + S_5 \right) \tag{15}$$

where

$$S_1 = \left(\frac{1}{\beta} + l \right) \sum_{q=0}^{p-1} \left(\frac{1}{\alpha} \right)^q \left(\frac{1}{\beta} \right)^{p-q+m-l+1}$$

$$S_2 = \sum_{q=0}^{p-1} \sum_{w=1}^{k-1} \left(\sum_{a=1}^{w} A_1 \left(\frac{1}{\alpha} \right)^{q+a} \left(\frac{1}{\beta} \right)^{m-l+1} + \sum_{b=1}^{p-q} A_2 \left(\frac{1}{\alpha} \right)^q \left(\frac{1}{\beta} \right)^{m+b-l+1} \right)$$

$$S_3 = l \sum_{q=0}^{p-1} \sum_{w=1}^{z+k-1} \left(\sum_{a=1}^{w} A_1 \left(\frac{1}{\alpha} \right)^{q+a} \left(\frac{1}{\beta} \right)^{m-l} + \sum_{b=1}^{p-q} A_2 \left(\frac{1}{\alpha} \right)^q \left(\frac{1}{\beta} \right)^{m+b-l} \right)$$

$$S_4 = \left(\frac{1}{\alpha}\right)^p \left(\frac{1}{\beta}\right)^{m-l+2} + \sum_{w=1}^{k-1} \left(\frac{1}{\alpha}\right)^{p+w} \left(\frac{1}{\beta}\right)^{m-l+1}$$

$$S_5 = l \left(\frac{1}{\alpha}\right)^p \left(\frac{1}{\beta}\right)^{m-l+1} + l \sum_{w=1}^{z+k-1} \left(\frac{1}{\alpha}\right)^{p+w} \left(\frac{1}{\beta}\right)^{m-l}$$

The equation (15) presents the final result of this article. In the special case (12) the equations (15) and (14) provide the same values, which supports the correctness of our result. Now, one can show that the headway distribution of the ASEP (after re-scaling) is of a Poissonian type for any parameters α, β (see Figure 1).

5 Conclusion

To conclude, the ASEP shows the Poissonian behavior for all possible values of the parameters α and β. Thus, this model can simulate only the traffic in free-flow regime, which means that

$$\lim_{s \to 0^+} P(s) \gg 0 \tag{16}$$

To be precise, we have to express that as the parameter β goes to zero ($\beta \to 0^+$) the Poissonian distribution changes since one can easily to follow that $P(s) = \delta(s) = \delta(k-1)$ for $\beta = 0$ and $\alpha \neq 0$. This holds true due to the shift of the probability $P(s) := P(k-1)$. The above mentioned shift is done because $P(k) = 0$ for every $k < 1$ in the original model. However, for every β we get the headway distribution satisfying the condition (16).

On contrary, in [7] a different variant of the ASE-model (high speed traffic model with open boundaries) is presented. The particles of this model also enter the system at the left and leave at the right of the chain. Moreover, all particles will move with their maximal possible speed, which is given by the speed limit v_{max} (a maximal length of the jump). For $v_{max} = 1$ one can obtain the model investigated in this article. Numerical simulations of the high speed model with $v_{max} \geq 2$ clearly show that the gap distribution can start from the origin (i.e. $\lim_{s \to 0^+} P(s) = 0$) for some values α, β.
Hence, the high speed model has the synchronized states as well, on contrary to the asymmetric simple exclusion process and it can be used to the full simulation of the real-road traffic.

6 Acknowledgement

Author is very grateful to V.B. Priezzhev and Petr Šeba for valuable advice and for the help with writing of this paper.

References

[1] B. Tilch and D. Helbing: Evaluation of single vehicle data in dependence of the vehicle-type, lane, and site, Traffic and Granular Flow '99, Springer-Verlag Berlin Heidelberg 2000, 333 – 338

[2] P. Seba and M. Krbalek: The statistical properties of the city transport in Cuernavaca (Mexico) and random matrix ensembles, J. Phys. A **33** (2000), L229 – L234

[3] M. Krbalek, P. Seba, and P. Wagner: Headways in traffic flow - remarks from a physical perspective, Phys. Rev. E. **64** (2001), 066119 – 066125

[4] B. Derrida: An exactly soluble non-equilibrium system: The asymmetric simple exclusion process, Physics Reports **301** (1998), 963 – 977

[5] B. Derrida, J. L. Lebowitz: Exact large deviation function in the asymmetric exclusion process, Phys. Rev. Lett. **80** (1998), 209 – 213

[6] N. Rajewski, L. Santen, A. Schadschneider, and M. Schreckenberg: The asymmetric exclusion process: Comparison of update procedures, Journal of Statistical Physics, **92** (1998), 151 – 194.

[7] J. de Gier: Exact stationary state for a deterministic high speed traffic model with open boundaries, J. Phys. A **34** (2001), 3707 – 3720

[8] M. Krbalek and P. Seba: Headway statistics of public transport in Mexican cities, to appear

7 Appendix

At the beginning, we cite some useful lemmas.

Lemma 1

Let m, n, a are the natural numbers and $a \leq m$. Then

$$\sum_{i=a}^{m} \left(\begin{array}{c} n-i+m-1 \\ n-1 \end{array} \right) = \left(\begin{array}{c} n+m-a \\ n \end{array} \right) \tag{17}$$

Proof: by means of the mathematical induction for the numbers m and n

Lemma 2

Assume that D, E are arbitrary matrices fulfilling the equation $DE = D + E$. Let m, n are the natural numbers. Then

$$D^n E^m = \sum_{i=1}^{m} \left(\begin{array}{c} n-i+m-1 \\ n-1 \end{array} \right) E^i + \sum_{j=1}^{n} \left(\begin{array}{c} n-j+m-1 \\ m-1 \end{array} \right) D^j \tag{18}$$

Proof: by means of the mathematical induction for the numbers m and n and (17)

Now we can rewrite the equation (11) in order to use the rules (5),(6). Thus, it is necessary to replace the matrices D on the right-hand side and matrices E to the left-hand side of the equation (11). The expression (8) yields

$$(D+E)^{N-i-k} = \sum_{m=1}^{N-i-k} \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!} \sum_{l=0}^{m} E^l D^{m-l}$$

Thus,

$$D\,(D+E)^{N-i-k} = \sum_{m=1}^{N-i-k} \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!} \sum_{l=0}^{m} D\,E^l D^{m-l} =$$

$$= \left| \begin{array}{c} D\,E^l = D + \sum_{z=1}^{l} E^z \\ D\,E^l D^{m-l} = D^{m-l+1} + \sum_{z=1}^{l} E^z D^{m-l} \end{array} \right| =$$

$$= \sum_{m=1}^{N-i-k} \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!} \sum_{l=0}^{m} \left(D^{m-l+1} + \sum_{z=1}^{l} E^z D^{m-l} \right)$$

It can be verified that

$$E^{k-1} D\,(D+E)^{N-i-k} =$$

$$= \sum_{m=1}^{N-i-k} \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!} \sum_{l=0}^{m} \left(E^{k-1} D^{m-l+1} + \sum_{z=1}^{l} E^{z+k-1} D^{m-l} \right)$$

Furthermore,

$$D\,E^{k-1} D\,(D+E)^{N-i-k} = \left| \begin{array}{c} D\,E^{k-1} = D + \sum_{w=1}^{k-1} E^w \\ D\,E^{z+k-1} = D + \sum_{w=1}^{z+k-1} E^w \end{array} \right| =$$

$$= \sum_{m=1}^{N-i-k} \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!}$$

$$\sum_{l=0}^{m} \left(D^{m-l+2} + \sum_{w=1}^{k-1} E^w D^{m-l+1} + l\,D^{m-l+1} + l \sum_{w=1}^{z+k-1} E^w D^{m-l} \right) \tag{19}$$

Another use of the expression (8) gives

$$(D+E)^{i-1} = \sum_{p=1}^{i-1} \frac{p\,(2i-p-3)!}{(i-1)!(i-p-1)!} \sum_{q=0}^{p} E^q D^{p-q} \tag{20}$$

Let now

$$\gamma = \gamma(m,k,i,N) = \frac{m\,(2N-2i-2k-m-1)!}{(N-i-k)!(N-i-k-m)!} \tag{21}$$

$$\delta = \delta(i,p) = \frac{p\,(2i-p-3)!}{(i-1)!(i-p-1)!} \tag{22}$$

Applying (21),(22), you can transform the equations (19) and (20) to

$$D\,E^{k-1}D\,(D+E)^{N-i-k} = \\ = \sum_{m=1}^{N-i-k}\sum_{l=0}^{m}\gamma\left(D^{m-l+2}+\sum_{w=1}^{k-1}E^{w}D^{m-l+1}+l\,D^{m-l+1}+\,l\sum_{w=1}^{z+k-1}E^{w}D^{m-l}\right)$$

$$(D+E)^{i-1}=\sum_{p=1}^{i-1}\sum_{q=0}^{p}\delta\,E^{q}D^{p-q}$$

Hence,

$$(D+E)^{i-1}D\,E^{k-1}D\,(D+E)^{N-i-k}=\sum_{p=1}^{i-1}\sum_{q=0}^{p}\sum_{m=1}^{N-i-k}\sum_{l=0}^{m}\delta\,\gamma \\ \left(E^{q}D^{p-q+m-l+2}+\sum_{w=1}^{k-1}E^{q}\,\overbrace{D^{p-q}E^{w}}\,D^{m-l+1}+\right. \\ \left. l\,E^{q}D^{p-q+m-l+1}\,+\,l\sum_{w=1}^{z+k-1}E^{q}\,\overbrace{D^{p-q}E^{w}}\,D^{m-l}\right) \tag{23}$$

In the expression (23) we can change the products indicated by the horizontal braces. Using (18) one can obtain

$$D^{p-q}E^{w}=\sum_{a=1}^{w}\binom{p-q-a+w-1}{p-q-1}E^{a}+\sum_{b=1}^{p-q}\binom{p-q-b+w-1}{w-1}D^{b} \tag{24}$$

Note that (24) is valid only for $p>q$. The remaining arrangement is the substitution of the (24) into the equation (23). With the help of the pre-factors

$$A_1=\binom{p-q-a+w-1}{p-q-1}$$

$$A_2=\binom{p-q-b+w-1}{w-1}$$

the expression (23) turns into the form

$$(D+E)^{i-1}D\,E^{k-1}D\,(D+E)^{N-i-k}=\sum_{p=1}^{i-1}\sum_{q=0}^{p-1}\sum_{m=1}^{N-i-k}\sum_{l=0}^{m}\delta\,\gamma \\ \left(E^{q}D^{p-q+m-l+2}+\,l\,E^{q}D^{p-q+m-l+1}+\sum_{w=1}^{k-1}\sum_{a=1}^{w}A_1E^{q+a}D^{m-l+1}\;+\right. \\ \left.+\sum_{w=1}^{k-1}\sum_{b=1}^{p-q}A_2E^{q}D^{m+b-l+1}\,+\,l\sum_{w=1}^{z+k-1}\sum_{a=1}^{w}A_1E^{q+a}D^{m-l}\,+\,l\sum_{w=1}^{z+k-1}\sum_{b=1}^{p-q}A_2E^{q}D^{b+m-l}\right)+ \\ +\sum_{p=1}^{i-1}\sum_{m=1}^{N-i-k}\sum_{l=0}^{m}\delta\,\gamma\left(E^{p}D^{m-l+2}+\sum_{w=1}^{k-1}E^{p+w}D^{m-l+1}+l\,E^{p}D^{m-l+1}+\,l\sum_{w=1}^{z+k+1}E^{p+w}D^{m-l}\right)$$

which together with the rules (5),(6) yields

$$\begin{aligned}
P^{(i)}(k) = \frac{\mathcal{A}}{Z_N} \sum_{p=1}^{i-1} \sum_{q=0}^{p-1} \sum_{m=1}^{N-i-k} \sum_{l=0}^{m} \delta\,\gamma \Bigg\{ & \left(\frac{1}{\alpha}\right)^{q} \left(\frac{1}{\beta}\right)^{p-q+m-l+2} + \, l \, \left(\frac{1}{\alpha}\right)^{q} \left(\frac{1}{\beta}\right)^{p-q+m-l+1} + \\
& + \sum_{w=1}^{k-1} \sum_{a=1}^{w} A_1 \left(\frac{1}{\alpha}\right)^{q+a} \left(\frac{1}{\beta}\right)^{m-l+1} + \sum_{w=1}^{k-1} \sum_{b=1}^{p-q} A_2 \left(\frac{1}{\alpha}\right)^{q} \left(\frac{1}{\beta}\right)^{m+b-l+1} + \\
& + \, l \sum_{w=1}^{z+k-1} \sum_{a=1}^{w} A_1 \left(\frac{1}{\alpha}\right)^{q+a} \left(\frac{1}{\beta}\right)^{m-l} + \, l \sum_{w=1}^{z+k-1} \sum_{b=1}^{p-q} A_2 \left(\frac{1}{\alpha}\right)^{q} \left(\frac{1}{\beta}\right)^{b+m-l} \Bigg\} + \\
& + \frac{\mathcal{A}}{Z_N} \sum_{p=1}^{i-1} \sum_{m=1}^{N-i-k} \sum_{l=0}^{m} \delta\,\gamma \Bigg\{ \left(\frac{1}{\alpha}\right)^{p} \left(\frac{1}{\beta}\right)^{m-l+2} + \sum_{w=1}^{k-1} \left(\frac{1}{\alpha}\right)^{p+w} \left(\frac{1}{\beta}\right)^{m-l+1} + \\
& + l \left(\frac{1}{\alpha}\right)^{p} \left(\frac{1}{\beta}\right)^{m-l+1} + \, l \sum_{w=1}^{z+k+1} \left(\frac{1}{\alpha}\right)^{p+w} \left(\frac{1}{\beta}\right)^{m-l} \Bigg\} \qquad (25)
\end{aligned}$$